

Table 1. Cotton variety performance in central Alabama, Macon Co. 2010*.						
Producers:	Mark & Ron Taylor					
ACES Cooperators:	L. Kuykendall, J. Clary, D. Monks, and J. Brasher					
Variety	Lint turnout	Lint yield (lbs/acre)	Mic	Length (inches)	Strength (g/tex)	Uniformity (%)
ST 5458B2RF	0.42	808	5.1	35.2	28.4	83.1
DP 1034B2RF	0.46	792	4.9	35.8	27.3	83.4
ST 5288B2RF	0.43	789	5.0	35.2	27.4	83.5
PHY 565WRF	0.42	782	4.8	36.5	30.9	84.9
DP 1048B2RF	0.45	767	4.6	36.5	27.7	83.0
ST 4288B2RF	0.38	750	4.7	36.5	28.9	83.3
DP 1050B2RF	0.45	747	4.7	36.2	27.3	83.6
PHY 375WRF	0.45	738	4.6	34.9	27.6	83.9
DP 0949B2RF	0.44	672	5.1	34.2	27.9	83.1
PHY 525RF	0.42	665	4.3	37.4	31.4	84.9
DP 555BR	0.44	647	4.6	35.2	29.5	83.1
PHY 485WRF	0.42	634	4.8	34.6	28.4	84.3
PHY 569WRF	0.42	593	4.9	34.6	29.4	83.9
*Note: Plots were 8, 40 inch rows wide X approximately 1600 ft. in length planted as single strips. Cotton was planted on May 11; and harvested on October 7, 2010.						
<i>Funding from Cotton Incorporated and the Alabama Cotton Commission was used in conducting this trial. We appreciate the support that Alabama cotton producers provide to our educational programs each year.</i>						