

<b>Table 4. Conventional cotton variety performance in central Alabama, Elmore Co. 2010*.</b>						
<b>Producer:</b>	Sanford Peeples					
<b>ACES Cooperators:</b>	L. Kuykendall, D. Monks, D. Delaney, K. Glass, and J. Brasher					
<b>Variety</b>	<b>Lint turnout</b>	<b>Lint yield (lbs/acre)</b>	<b>Mic</b>	<b>Length (32nd)</b>	<b>Strength (g/tex)</b>	<b>Uniformity (%)</b>
CT LINWOOD	0.41	719	4.4	33.9	28.4	82.2
GA EX 230	0.41	690	4.6	32.3	25.9	80.7
DP 555BR	0.42	682	4.9	32.6	24.7	80.9
LSU 1110017	0.39	657	4.2	36.2	26.8	78.9
PHY 440W	0.40	630	3.9	35.2	29.6	82.8
X1-90	0.39	611	4.1	34.9	28.5	81.7
CT 212	0.39	603	3.9	33.6	28.0	79.9
CT 310	0.37	596	4.0	34.6	28.0	81.8
CT 210	0.39	576	4.4	33.6	27.1	80.8
FM 1845LLB2	0.37	575	3.9	36.2	31.1	82.1
DP 491	0.40	539	3.9	34.6	29.8	82.8
*Note: Plots were 4, 38 inch rows wide X approximately 1000 ft. in length no-till planted with 3 strips per variety. Planting date was May 6; harvest date was September 21, 2010.						
<i>Funding from Cotton Incorporated and the Alabama Cotton Commission was used in conducting this trial. We appreciate the support that Alabama cotton producers provide to our educational programs each year.</i>						